arXiv:1508.07315v1 [math.CO] 28 Aug 2015

On linear equations arising in Combinatorics (Part II)

Masood Aryapoor*

Abstract

In [1], the author introduces the class of Farkas–related vectors for which a version of Farkas' lemma over integers is derived. In this paper, being the second part of [1], two similar classes are introduced and studied. The terminology and results of [1] are freely used in this article.

1 Almost Farkas–related vectors

We begin with the following definition, compare with the definition of Farkas–related vectors in [1].

Definition 1.1. *Vectors* $v_1, ..., v_m \in \mathbb{Z}^n$ *are called almost Farkas–related or a.f.r for short, if the following condition holds: Let* $a_1 \leq b_1, ..., a_m \leq b_m$ *be arbitrary integers. If a vector* $w \in \sum_{a_i=b_i} a_i v_i + \sum_{a_i \neq b_i} \mathbb{Z} v_i$ *can be written as* $w = \sum_{i=1}^{m} x_i v_i$ *for rational numbers* $a_1 \leq x_1 \leq b_1, ..., a_m \leq x_m \leq b_m$*, then there exist integers* $a_1 \leq y_1 \leq b_1, ..., a_m \leq y_m \leq b_m$ *such that* $w = \sum_{i=1}^{m} y_i v_i$.

1.1 Characterizations of almost Farkas–related vectors

We want to obtain simple and useful characterizations of almost Farkas–related vectors. First we present the following characterization.

Lemma 1.1. *Let* $v_1, ..., v_m \in \mathbb{Z}^n$ *be arbitrary vectors. Then the following conditions are equivalent.*
(1) The vectors $v_1, ..., v_m \in \mathbb{Z}^n$ *are a.f.r.*
(2) Let $I \subset \{1, ..., m\}$ *be an arbitrary nonempty set. For arbitrary integers* $a_i < b_i$ $(i \in I)$*, if a vector* $w \in \sum_{i \in I} \mathbb{Z} v_i$ *can be written as* $w = \sum_{i \in I} x_i v_i$ *for rational numbers* $a_i \leq x_i \leq b_i$ $(i \in I)$*, then there exist integers* $a_i \leq y_i \leq b_i$ $(i \in I)$ *such that* $w = \sum_{i \in I} y_i v_i$.
(3) Let $I \subset \{1, ..., m\}$ *be an arbitrary nonempty set. If a vector* $w \in \sum_{i \in I} \mathbb{Z} v_i$ *can be written as* $w = \sum_{i \in I} x_i v_i$ *for rational numbers* $0 \leq x_i \leq 1$ $(i \in I)$*, then there exist numbers* $y_i \in \{0, 1\}$ $(i \in I)$ *such that* $w = \sum_{i \in I} y_i v_i$.

---

*E-mail: aryapoor2002@yahoo.com

(4) Let $I \subset \{1, ..., m\}$ be an arbitrary nonempty set and k be a natural number. If a vector $w \in \sum_{i\in I} \mathbb{Z}v_i$ can be written as $kw = \sum_{i\in I} a_i v_i$ for integers $0 \le a_i \le k$ $(i \in I)$, then there exist numbers $y_i \in \{0, 1\}$ $(i \in I)$ such that $w = \sum_{i\in I} y_i v_i$.

Proof. It is straightforward to check that (1)⇔(2) and (3)⇔(4) . Clearly we have (2)⇒(3). Therefore it remains to prove (3)⇒(2).

Suppose that (3) holds. Let $I \subset \{1, ..., m\}$ be a nonempty set and $a_i < b_i$ $(i \in I)$ be integers. Suppose that a vector $w \in \sum_{i\in I} \mathbb{Z}v_i$ can be written as $w = \sum_{i\in I} x_i v_i$ for rational numbers $a_i \le x_i \le b_i$ $(i \in I)$. Set $J = \{i \in I | x_i \ne b_i\}$. We have

$$w - \sum_{i\in I, i\notin J} (b_i - 1)v_i - \sum_{i\in J} [x_i]v_i = \sum_{i\in J} (x_i - [x_i])v_i + \sum_{i\in I, i\notin J} v_i \in \sum_{i\in I} \mathbb{Z}v_i.$$

So, by (3), there exist numbers $y_i \in \{0, 1\}$ such that

$$w - \sum_{i\in I, i\notin J} (b_i - 1)v_i - \sum_{i\in J} [x_i]v_i = \sum_{i\in I} y_i v_i.$$

This equality can be rewritten as $w = \sum_{i\in I, i\notin J}(b_i - 1 + y_i)v_i + \sum_{i\in J}([x_i] + y_i)v_i$. Note that the coefficients are integers. Since $a_i < b_i$, we have $a_i \le b_i - 1 + y_i \le b_i$ for $i \notin J$. Moreover, for $i \in J$, we have $a_i \le [x_i] \le b_i - 1$ which implies that $a_i \le [x_i] + y_i \le b_i$ for $i \in J$ as well. The proof of (3)⇒(2) is therefore complete. □

A useful criterion is given below, consult [1] to see the definition of "elementary integral relations".

Proposition 1.2. *Vectors $v_1, ..., v_m \in \mathbb{Z}^n$ are a.f.r if and only if for every elementary integral relation $\sum_{i=1}^m a_i v_i = 0$, we have $|a_1|, ..., |a_m| \le 2$ and moreover there exists at most one $1 \le i \le m$ such that $|a_i| = 2$.*

Proof. First, the "only if direction" is proved. Suppose that $v_1, ..., v_m$ are a.f.r and let $\sum_{i=1}^m a_i v_i = 0$ be an elementary integral relation. Without loss of generality, we may assume that $\{i | a_i \ne 0\} = \{1, ..., r\}$. By symmetry, it is enough to show that $|a_1| \in \{1, 2\}$ and if $|a_1| = 2$ then $a_2, ..., a_r \in \{-1, 1\}$. Since $0 = v_1 + \frac{a_2}{a_1}v_2 + \cdots + \frac{a_r}{a_1}v_r$, using the fact that $v_1, ..., v_m$ are a.f.r, we see that there exist integers

$$1 \le y_1 \le 2, [\frac{a_2}{a_1}] \le y_2 \le [\frac{a_2}{a_1}] + 1, ..., [\frac{a_r}{a_1}] \le y_r \le [\frac{a_r}{a_1}] + 1,$$

such that $0 = y_1 v_1 + \cdots + y_r v_r$. It is easy to see that the set of vectors $(x_1, ..., x_r) \in \mathbb{Z}^r$ satisfying $\sum_{i=1}^r x_i v_i = 0$, is equal to $\mathbb{Z}(a_1, ..., a_r)$. It follows that $(y_1, ..., y_r) = l(a_1, ..., a_r)$ for an integer l. Since $y_1 = 1$ or $y_2 = 2$, we conclude that $|a_1| \in \{1, 2\}$. Now suppose that $a_1 = 2$. Then $(y_1, ..., y_r) = (a_1, ..., a_r)$, i.e. $[\frac{a_i}{2}] \leq y_i = a_i \leq [\frac{a_i}{2}] + 1$ for $i = 2, ..., r$. But in this case, there also exist integers

$$0 \leq z_1 \leq 1, [\frac{a_2}{2}] \leq z_2 \leq [\frac{a_2}{2}] + 1, ..., [\frac{a_r}{2}] \leq z_r \leq [\frac{a_r}{2}] + 1,$$

such that $0 = z_1v_1 + \cdots + z_rv_r$. We have $(z_1, ..., z_r) \in \mathbb{Z}(2, a_2, ..., a_r)$ and $z_1 = 0$ or $z_1 = 1$. It follows that $z_1 = 0$ and consequently $z_2 = \cdots = z_r = 0$ which implies that $[\frac{a_i}{2}] \in \{-1, 0\}$ where $i = 2, ..., r$. This means that $a_i \in \{-2, -1, 1\}$ where $i = 2, ..., r$. But, since $[\frac{a_i}{2}] \leq a_i \leq [\frac{a_i}{2}] + 1$ for every $i = 2, ..., r$, it is easy to see that we must have $a_i \in \{-1, 1\}$ for every $i = 2, ..., r$. If $a_1 = -2$, then one uses $(-a_1)v_1 + \cdots + (-a_r)v_r = 0$ to show that $-a_i \in \{-1, 1\}$ for every $i = 2, ..., r$ which is obviously the same as $a_i \in \{-1, 1\}$ for every $i = 2, ..., r$.

Now we prove the "if direction". First note that without loss of generality we may assume that none of the vectors $v_1, ..., v_m$ are zero. We use induction on m to prove that the vectors $v_1, ..., v_m$ satisfy Part (4) of Lemma 1.1. So let $I \subset \{1, ..., m\}$ be a nonempty set and suppose that a vector $w \in \sum_{i \in I} \mathbb{Z}v_i$ can be written as $kw = \sum_{i \in I} a_iv_i$ for integers $0 \leq a_i \leq k$ $(i \in I)$, where k is an arbitrary natural number. We need to show that the vector w can be written as $w = \sum_{i \in I} y_iv_i$ where $y_i \in \{0, 1\}$ $(i \in I)$. First let $m = 1$. Then we must have $I = \{1\}$ and since $w = av_1$ for an integer a, we have $a_1v_1 = kw = kav_1$ or $(a_1 - ka)v_1 = 0$. If $w = 0$, then $w = 0.v_1$ and we are done. Otherwise, we must have $a = 1$ because $0 \leq a_1 \leq k$ and $a_1 - ka = 0$. So $w = v_1$ and we are done with the case $m = 1$.

Next, we prove the inductive step. If $I \neq \{1, ..., m\}$, then we are done, because the vectors v_i $(i \in I)$ obviously satisfy the relevant condition on their elementary integral relations and therefore are almost Farkas–related by induction. So we may assume that $I = \{1, ..., m\}$. I claim that there exists $J \subset \{1, ..., m\}$ such that the vectors $v_i (i \in J)$ form a basis for $\sum_{i=1}^m \mathbb{Z}v_i$ over $\mathbb{Z}$. To prove this, take a set $J \subset \{1, ..., m\}$ such that $v_i (i \in J)$ are linearly independent and there does not exist a subset $J' \subset \{1, ..., m\}$ with the property that the vectors $v_i (i \in J')$ are linearly independent and $\sum_{i \in J} \mathbb{Z}v_i \subsetneq \sum_{i \in J'} \mathbb{Z}v_i$. If $v_j \notin \sum_{i \in J} \mathbb{Z}v_i$ for an index $1 \leq j \leq m$, then the vectors $v_j, v_i (i \in J)$ are linearly dependent since otherwise $\sum_{i \in J} \mathbb{Z}v_i \subsetneq \sum_{i \in J \cup \{j\}} \mathbb{Z}v_i$ would contradict the choice of J. It follows that there exists an elementary integral relation between $v_j, v_i (i \in J)$. This relation has to be of the form $2v_j + \sum_{i \in J} a_iv_i = 0$ where $a_i \in \{-1, 0, 1\}$ for every $i \in I$. Choose $i \in J$ such that $a_i \neq 0$ and set $J' = \{j\} \cup (J \setminus \{i\})$. Then the vectors $v_i (v_i \in J')$ are linearly independent and moreover $\sum_{i \in J} \mathbb{Z}v_i \subsetneq \sum_{i \in J'} \mathbb{Z}v_i$, a contradiction. It follows that the vectors $v_i (i \in J)$ generate $\sum_{i=1}^m \mathbb{Z}v_i$ and the claim is proved. Without loss of generality, we may assume that $J = \{v_1, ..., v_r\}$.

If $r = m$, then the identity $w = \sum_{i=1}^m \frac{a_i}{k} v_i$ and the fact $w \in \sum_{i=1}^m \mathbb{Z}v_i$ imply that $\frac{a_1}{k}, ..., \frac{a_m}{k} \in \mathbb{Z}$. But $0 \leq a_1, ..., a_m \leq k$, so we must have $a_i \in \{0, k\}$, and consequently $w = \sum_{i=1}^m \frac{a_i}{k} v_i$ is the desired presentation. In what follows, we may therefore assume that $r < m$.

First suppose that there does not exist $1 \leq i \leq r$ such that $v_i \in \sum_{i \neq j=1}^m \mathbb{Z}v_j$. For every $r < j \leq m$, we have $v_j \in \sum_{i=1}^r \mathbb{Z}v_i$, i.e. $\sum_{i=1}^r b_iv_i + v_j = 0$ for some

$b_1, ..., b_r \in \mathbb{Z}v_j$. It is easy to see that this relation is in fact an elementary integral relation. So there is at most one b_i with $|b_i| > 1$. If there is some b_i with $|b_i| = 1$ then we clearly have $v_i \in \sum_{i \neq j=1}^{m} \mathbb{Z}v_j$. It follows that exactly one b_i is not zero and it must be equal to ± 2, since otherwise $v_i \in \sum_{i \neq j=1}^{m} \mathbb{Z}v_j$. It follows from this discussion that the set $\{1, ..., m\}$ can be partitioned into r disjoint subsets $I_1, ..., I_r$ such that for every $i = 1, ..., r$, we have $i \in I_i$ and $v_j = \pm 2v_i$ for every $j \in I_i$. In this case, if $w = d_1v_1 + \cdots + d_rv_r$, then $k(d_iv_i) = \sum_{l \in I_i} a_lv_l$. Now, if $r > 1$, then we can use induction, to find suitable presentations for each d_iv_i and by adding them together, one can find an acceptable representation for w. So we may assume that $r = 1$. To proceed, we set $I_{\pm} = \{2 \leq i \leq m | v_i = \pm 2v_1\}$. We can write $kw = (a_1 + 2\sum_{i \in I_+} a_i - 2\sum_{i \in I_-} a_i)v_1$. On the other hand we have $w = dv_1$ where $d \in \mathbb{Z}$. So $kd = a_1 + 2\sum_{i \in I_+} a_i - 2\sum_{i \in I_-} a_i$. Since

$$-2|I_-|k \leq a_1 + 2\sum_{i \in I_+} a_i - 2\sum_{i \in I_-} a_i \leq k + 2|I_+|k,$$

we conclude that $-2|I_-| \leq d \leq 1 + 2|I_+|$. Using this inequality, one can see that there are integers $0 \leq l_1 \leq |I_+|, 0 \leq l_2 \leq |I_-|$ and $l_3 \in \{0, 1\}$ such that $d = l_3 + 2l_1 - 2l_2$. By taking arbitrary sets $L_1 \subset I_+$ and $L_2 \subset I_2$ satisfying $|L_1| = l_1$ and $|L_2| = l_2$, we obtain $w = l_3v_1 + \sum_{i \in L_1 \cup L_2} v_i$ which is the desired presentation and we are done. So we may assume that there exists $1 \leq i \leq r$ such that $v_i \in \sum_{i \neq j=1}^{m} \mathbb{Z}v_j$. Without loss of generality we may assume that $v_1 \in \sum_{j=2}^{m} \mathbb{Z}v_j$.

It is easy to see that for every vector $u \in \sum_{i=1}^{m} \mathbb{Z}v_i$, there is a unique vector $p(u) \in \sum_{i=2}^{r} \mathbb{Z}v_i$, such that $u - p(u) \in \mathbb{Z}v_1$. Moreover the map $p : \sum_{i=1}^{m} \mathbb{Z}v_i \to \sum_{i=2}^{r} \mathbb{Z}v_i$ is a $\mathbb{Z}$–linear map whose kernel is $\mathbb{Z}v_1$. I claim that the vectors $p(v_2), ..., p(v_m)$, satisfy the condition in the proposition. In fact, if $\sum_{i=2}^{m} b_ip(v_i) = 0$ is an elementary integral relation, then we have $\sum_{i=2}^{m} b_iv_i = -b_1v_1$ for an integer b_1. It is easy to see that the relation $\sum_{i=1}^{m} b_iv_i = 0$ is an elementary integral relation. So $b_1, b_2, ..., b_m$ satisfy the relevant condition which implies that $b_2, ..., b_m$ satisfy the condition as well. This proves the claim. In particular, by induction, we conclude that the vectors $p(v_2), ..., p(v_m)$ are a.f.r. Since $w \in \sum_{i=1}^{m} \mathbb{Z}v_i$, we have $p(w) \in \sum_{i=2}^{m} \mathbb{Z}p(v_i)$. Moreover, we have $kp(w) = \sum_{i=2}^{m} a_ip(v_i)$. It follows that there exist numbers $y_2, ..., y_m \in \{0, 1\}$ such that $p(w) = \sum_{i=2}^{m} y_ip(v_i)$. Since the kernel of p is $\mathbb{Z}v_1$, we conclude that $w = av_1 + \sum_{i=2}^{m} y_iv_i$ for an integer a. If $a_1 - ak = 0$, then we must have $a_1 = a = 0$ or $a_1 = k, a = 1$ because $0 \leq a_1 \leq k$. In both cases, we obtain an acceptable presentation for w, and we are done. So suppose that $a_1 - ak \neq 0$. At this stage, we consider two cases.

Case 1: $a < 0$. Then

$$(a_1 - ak)w = (a_1 - ak - k)(av_1 + \sum_{i=2}^{m} y_iv_i) + \sum_{i=1}^{m} a_iv_i =$$

$$= (a+1)(a_1 - ak)v_1 + \sum_{i=2}^{m}((a_1 - ak - k)y_i + a_i)v_i.$$

So we have $(a_1 - ak)(w - (a+1)v_1) = \sum_{i=2}^{m}((a_1 - ak - k)y_i + a_i)v_i$. Note that we have $a_1 - ak > 0$. It is easy to see that $0 \leq (a_1 - ak - k)y_i + a_i \leq a_1 - ak$ for each $i = 2, ..., m$. Since $w - (a+1)v_1 \in \sum_{i=2}^{m} \mathbb{Z}v_i$, by induction there exist numbers $y'_2, ..., y'_m \in \{0, 1\}$, such that $w = (a+1)v_1 + y'_2v_2 + \cdots + y'_mv_m$. Continuing this process, since a is negative, will produce a presentation $w = \sum_{i=2}^{m} z_iv_i$ where $z_2, ..., z_m \in \{0, 1\}$, and hence we are done.

Case 2: $a \geq 0$. In this case if $a \leq 1$, then we are done. Otherwise, the proof is similar to the previous case. One just uses the following identity,

$$(ak - a_1)(w - (a-1)v_1) = \sum_{i=2}^{m}((ak - a_1 - k)y_i + a_i)v_i.$$

□

1.2 Farkas' Lemma for almost Farkas–related vectors

Using the definition of a.f.r vectors, one can derive the following theorem, consult [1].

Theorem 1.3. *Suppose that vectors $v_1, ..., v_m \in \mathbb{Z}^n$ are almost Farkas–related and let arbitrary integers $a_1 \leq b_1, ..., a_m \leq b_m$ be given. Then a vector $w \in \mathbb{Z}^n$ can be written as $w = \sum_{i=1}^{m} x_iv_i$ where $a_1 \leq x_1 \leq b_1, ..., a_m \leq x_m \leq b_m$ are integers if and only if $w - \sum_{a_i = b_i} a_iv_i \in \sum_{a_i \neq b_i} \mathbb{Z}v_i$ and*

$$(u, w) \leq \sum_{i=1}^{m} a_i \frac{(u, v_i) - |(u, v_i)|}{2} + \sum_{i=1}^{m} b_i \frac{(u, v_i) + |(u, v_i)|}{2},$$

for every $\{v_1, ..., v_m\}$–indecomposable point $[u] \in \mathbb{RP}_+^{n-1}$.

1.3 Almost Farkas Graphs

We follow the notations in Section 4.1 of [1].

Definition 1.2. *A connected (simple) graph G is called almost Farkas if the vectors $\{v(e)\}_{e \in E(G)}$ are almost Farkas-related.*

Here is a characterization of almost Farkas graphs.

Theorem 1.4. *A connected graph G is almost Farkas if and only if every two disjoint simple odd cycles of G cover the whole graph, i.e every vertex of G belongs to one of the cycles.*

Proof. First suppose that G is almost Farkas. Assume on the contrary that there exist two disjoint simple odd cycles of G that do not cover the whole graph. It follows that there exist a simple path of length > 1 which connects the two cycles. This gives rise to an elementary integral relation having at least two coefficients equal to 2 or -2 (see Proposition 4.2 in [1]), a contradiction by Proposition 1.2.

To prove the other direction, we use Proposition 1.2. An elementary integral relation of G that violates the condition in Proposition 1.2 correspond to two disjoint odd cycles of G that are connected by a path of length > 1, see Proposition 4.2 in [1]. This is however not possible since otherwise such two odd cycles would not cover the whole graph. □

The graph, drawn below, is an almost Farkas graph. But this graph is not a Farkas graph, see [1].



2 Weakly Farkas–related vectors

The definition of weakly Farkas–related vectors is given below.

Definition 2.1. *Vectors $v_1, ..., v_m \in \mathbb{Z}^n$ are said to be $\underline{\text{weakly}}$ Farkas–related, or w.f.r for short, if the following condition holds: For arbitrary integers $a_1 < b_1, ..., a_m < b_m$, if a vector $w \in \sum_{i=1}^m \mathbb{Z}v_i$ can be written as $w = \sum_{i=1}^m x_i v_i$ for rational numbers $a_1 \leq x_1 \leq b_1, ..., a_m \leq x_m \leq b_m$, then there exist integers $a_1 \leq y_1 \leq b_1, ..., a_m \leq y_m \leq b_m$ such that $w = \sum_{i=1}^m y_i v_i$.*

2.1 A characterization of weakly Farkas–related vectors

We begin with the following lemma.

Lemma 2.1. *Let $v_1, ..., v_m \in \mathbb{Z}^n$ be arbitrary vectors. Then the following conditions are equivalent.*
(1) The vectors $v_1, ..., v_m \in \mathbb{Z}^n$ are w.f.r.
(2) For arbitrary integers $a_1 < b_1, ..., a_m < b_m$, if there exist rational numbers $a_1 \leq x_1 \leq b_1, ..., a_m \leq x_m \leq b_m$ such that $\sum_{i=1}^m x_i v_i = 0$, then there exist integers $a_1 \leq y_1 \leq b_1, ..., a_m \leq y_m \leq b_m$ such that $\sum_{i=1}^m y_i v_i = 0$.
(3) For arbitrary integers $a_1, ..., a_m$, if there exist rational numbers $a_1 \leq x_1 \leq a_1 + 1, ..., a_m \leq x_m \leq a_m + 1$ such that $\sum_{i=1}^m x_i v_i = 0$, then there exist integers $a_1 \leq y_1 \leq a_1 + 1, ..., a_m \leq y_m \leq a_m + 1$ such that $\sum_{i=1}^m y_i v_i = 0$.

Proof. It is obvious that (1)⇒(2) and (2)⇒(3). To prove (2)⇒(1), suppose that integers $a_1 < b_1, ..., a_m < b_m$ are given and a vector $w \in \sum_{i=1}^m \mathbb{Z}v_i$ can be written as $w = \sum_{i=1}^m x_i v_i$ for rational numbers $a_1 \leq x_1 \leq b_1, ..., a_m \leq x_m \leq b_m$. Since $w \in \sum_{i=1}^m \mathbb{Z}v_i$, there exist $c_1, ..., c_m \in \mathbb{Z}$ such that $w = \sum_{i=1}^m c_i v_i$. We have $\sum_{i=1}^m (x_i - c_i)v_i = 0$, where

$$a_1 - c_1 \leq x_1 - c_1 \leq b_1 - c_1, ..., a_m - c_m \leq x_m - c_m \leq b_m - c_m.$$

Therefore there exist integers

$$a_1 - c_1 \leq y_1 \leq b_1 - c_1, ..., a_m - c_m \leq y_m \leq b_m - c_m$$

such that $\sum_{i=1}^m y_i v_i = 0$. The presentation $w = \sum_{i=1}^m (y_i + c_i)v_i$ is the desired presentation because

$$a_1 \leq y_1 + c_1 \leq b_1, ..., a_m \leq y_m + c_m \leq b_m.$$

To prove (3)⇒(2), suppose that integers $a_1 < b_1, ..., a_m < b_m$ are given and $\sum_{i=1}^m x_i v_i = 0$ for rational numbers $a_1 \leq x_1 \leq b_1, ..., a_m \leq x_m \leq b_m$. Setting $c_i = [x_i]$ if $x_i \neq b_i$ and $c_i = [x_i] - 1$ if $x_i = b_i$, we have

$$c_1 \leq x_1 \leq c_1 + 1, ..., c_m \leq x_m \leq c_m + 1.$$

Therefore there exist integers

$$c_1 \leq y_1 \leq c_1 + 1, ..., c_m \leq y_m \leq c_m + 1,$$

such that $\sum_{i=1}^m y_i v_i = 0$. We clearly have $a_1 \leq y_1 \leq b_1, ..., a_m \leq y_m \leq b_m$ and we are done.

□

A relatively simple characterization of w.f.r vectors is given in the following proposition.

Proposition 2.2. *Let $v_1, ..., v_m \in \mathbb{Z}^n$ be arbitrary vectors. The vectors $v_1, ..., v_m$ are w.f.r if and only if the following condition holds: Suppose that integers $a_1, ..., a_m \in \{-1, 0, 1\}$ are given such that there exists only one index $1 \leq s \leq m$ with $a_s = 1$. If there exist rational numbers*

$$a_1 \leq x_1 \leq a_1 + 1, ..., a_m \leq x_m \leq a_m + 1$$

such that $\sum_{i=1}^m x_i v_i = 0$, then there exist integers

$$a_1 \leq y_1 \leq a_1 + 1, ..., a_m \leq y_m \leq a_m + 1$$

such that $\sum_{i=1}^m y_i v_i = 0$.

Proof. Using Lemma 2.1, one only needs to prove that the desired property is equivalent to Property (3) in Lemma 2.1. Clearly if Property (3) holds in Lemma 2.1, then the property in the proposition holds as well.

To prove the converse, suppose that integers $a_1, ..., a_m$ are given and there exist rational numbers $a_1 \leq x_1 \leq a_1 + 1, ..., a_m \leq x_m \leq a_m + 1$ such that $\sum_{i=1}^{m} x_i v_i = 0$. We need to prove that there exist integers

$$a_1 \leq y_1 \leq a_1 + 1, ..., a_m \leq y_m \leq a_m + 1$$

such that $\sum_{i=1}^{m} y_i v_i = 0$. We prove this by induction on $\sum_{i=1}^{m} |2a_i + 1|$. If $\sum_{i=1}^{m} |2a_i + 1| \leq m$, then $a_1, ..., a_m \in \{-1, 0\}$ in which case the numbers $y_1 = ... = y_n = 0$ are the desired numbers.

Now we prove the inductive step. We consider three cases.

Case 1: There exists $1 \leq r \leq m$ such that $|x_r| = max_{i=1,...,m}|x_i|$ and $a_r > 0$. Without loss of generality we may assume $r = 1$. We define $a'_1, ..., a'_m$ as follows

$$a'_i = \begin{cases} 1 & i = 1 \\ 0 & a_i \geq 0 \text{ and } i > 1 \\ -1 & a_i < 0 \end{cases}$$

It is easy to see that $a'_i \leq \frac{x_i}{x_1} \leq a'_i + 1$ for every $i = 1, ..., m$. Since $\sum_{i=1}^{m} \frac{x_i}{x_1} v_i = 0$ and the numbers $a'_1, ..., a'_m$ satisfy the condition in the proposition, we conclude that there exist integers

$$a'_1 \leq y'_1 \leq a'_1 + 1, ..., a'_m \leq y'_m \leq a'_m + 1$$

such that $\sum_{i=1}^{m} y'_i v_i = 0$. Consider the numbers $a_1 - y'_1, ..., a_m - y'_m$. I claim that $\sum_{i=1}^{m} |2(a_i - y'_i) + 1| < \sum_{i=1}^{m} |2a_i + 1|$. In fact we have $|2(a_1 - y'_1) + 1| < |2a_1 + 1|$ because $y'_1 \in \{1, 2\}$ and $a_1 > 0$. So it is enough to show $|2(a_i - y'_i) + 1| \leq |2a_i + 1|$ for every $i = 2, ..., m$. If $a_i \geq 0$, then $y'_i \in \{0, 1\}$ and therefore $|2(a_i - y'_i) + 1| \leq |2a_i + 1|$ (note that $|2(0 - 1) + 1| = |2(0) + 1|$). Similarly if $a_i < 0$, then $y'_i \in \{-1, 0\}$ which implies that $|2(a_i - y'_i) + 1| \leq |2a_i + 1|$. Now, we have $\sum_{i=1}^{m} (x_i - y'_i) v_i = 0$ where

$$a_1 - y'_1 \leq x_1 - y'_1 \leq (a_1 - y'_1) + 1, ..., a_m - y'_m \leq x_m - y'_m \leq (a_m - y'_m) + 1.$$

Therefore, by induction, there exist integers

$$a_1 - y'_1 \leq y''_1 \leq (a_1 - y'_1) + 1, ..., a_m - y'_1 \leq y''_m \leq (a_m - y'_1) + 1,$$

such that $\sum_{i=1}^{m} y''_i v_i = 0$. Finally, we have $\sum_{i=1}^{m} (y'_i + y''_i) v_i = 0$ where

$$a_1 \leq y'_1 + y''_1 \leq a_1 + 1, ..., a_m \leq y'_m + y''_m \leq a_m + 1,$$

and we are done.

Case 2: For all $1 \leq i \leq m$ satisfying $|x_r| = max_{j=1,...,m}|x_j|$ we have $a_i \leq 0$, and there exists $1 \leq r \leq m$ such that $|x_r| = max_{j=1,...,m}|x_j|$ and $a_r = 0$. In this case, we have $|x_r| \leq 1$ which implies that $|x_i| \leq 1$ for every $i = 1, ..., m$. Since $a_i \leq x_i \leq a_{i+1}$ for every $i = 1, ..., m$, we must have $-1 \leq a_i \leq 1$ for every $i = 1, ..., m$. If there exists an index $1 \leq s \leq m$ such that $a_s = 1$, then we have $x_s = 1$, since $|x_s| \leq 1$ and $a_s = 1 \leq x_s \leq a_s + 1 = 2$. But

then $|x_s| = max_{j=1,...,m}|x_j|$ and $a_s > 0$, a contradiction. Therefore we have $a_i \in \{-1, 0\}$ for every $i = 1, ..., m$. It means that the numbers $a_1, ..., a_m$ satisfy the condition in the proposition and we are therefore done.
Case 3: For all $1 \leq i \leq m$ satisfying $|x_i| = max_{j=1,...,m}|x_j|$, we have $a_i < 0$. Set $b_1 = -a_1 - 1, ..., b_m = -a_m - 1$. We have

$$b_1 \leq -x_1 \leq b_1 + 1, ..., b_m \leq -x_m \leq b_m + 1.$$

Moreover for all $1 \leq i \leq m$ satisfying $| - x_i| = max_{i=1,...,m}| - x_i|$, we have $b_i \geq 0$. Since $\sum_{i=1}^{m} |2b_i + 1| = \sum_{i=1}^{m} |2a_i + 1|$ in view of the identity $|2a + 1| = |2(-a - 1) + 1|$, we may use Case 1 or Case 2 to obtain integers

$$b_1 \leq y_1 \leq b_1 + 1, ..., b_m \leq y_m \leq b_m + 1,$$

such that $\sum_{i=1}^{m} y_i v_i = 0$. The numbers $-y_1, ..., -y_m$ satisfy the desired condition, i.e.

$$a_1 \leq -y_1 \leq a_1 + 1, ..., a_m \leq -y_m \leq a_m + 1$$

and $\sum_{j=1}^{m}(-y_j)v_j = 0$. Therefore the proof is complete.

□

Compared with Proposition 2.1, it is substantially easier to use Proposition 2.2 in order to check if given vectors are w.f.r. It could however be possible to obtain a simpler characterization that is similar to the characterization of weakly Farkas-related vectors given in Proposition 1.2.

2.2 Farkas' Lemma for weakly Farkas–related vectors

Using the definition of w.f.r vectors, one can derive the following theorem.

Theorem 2.3. *Suppose that vectors $v_1, ..., v_m \in \mathbb{Z}^n$ are weakly Farkas–related and let arbitrary integers $a_1 < b_1, ..., a_m < b_m$ be given. Then a vector $w \in \mathbb{Z}^n$ can be written as $w = \sum_{i=1}^{m} x_i v_i$ where $a_1 \leq x_1 \leq b_1, ..., a_m \leq x_m \leq b_m$ are integers if and only if $w \in \sum_{i=1}^{m} \mathbb{Z}v_i$ and*

$$(u, w) \leq \sum_{i=1}^{m} a_i \frac{(u, v_i) - |(u, v_i)|}{2} + \sum_{i=1}^{m} b_i \frac{(u, v_i) + |(u, v_i)|}{2},$$

for every $\{v_1, ..., v_m\}$–indecomposable point $[u] \in \mathbb{RP}_+^{n-1}$.

2.3 Weakly Farkas Graphs

We follow the notations in Section 4.1 of [1].

Definition 2.2. *A connected simple graph G is called weakly Farkas if the vectors $\{v(e)\}_{e \in E(G)}$ are weakly Farkas-related.*

A graph is said to satisfy the odd-cycle condition if for every two vertex-disjoint simple odd cycles of the graph there exist two vertexes, one from each cycle, that are connected by an edge, also see [2].

Theorem 2.4. *A connected simple graph is weakly Farkas if and only if it satisfies the odd-cycle condition.*

Proof. To see a proof of the "if direction", see [2]. To prove the "only if direction", suppose that a graph G does not satisfy the odd-cycle condition. We need to show that G is not weakly Farkas. There exist two (simple) odd cycles $u_1u_2...u_m$ and $v_1v_2...v_n$ whose distance is greater than 1. Since G is connected, there is a path, say $u_1w_1...w_pv_1$, connecting these two cycles. We may assume that $\{w_1,...,w_p\} \cap \{u_1,...,u_m,v_1,...,v_n\} = \emptyset$. Among such cycles and paths joining them, we choose two cycles and a path joining them such that $m+n+p$ is as small as possible. The following properties hold:
(1) There are no edges between the vertexes $u_1, w_1, ..., w_p, v_1$ except the edges of the path $u_1w_1...w_pv_1$. In fact if there were such an edge then we would get a smaller path joining the two cycles, a contradiction.
(2) There are no edges between the vertexes $u_1, ..., u_m$ except the edges of the cycle $u_1u_2...u_m$. The same holds for the cycle $v_1v_2...v_n$. To see this, suppose that two vertexes v_i and v_j are connected by an edge where $1 \leq i < j-1 \leq n-1$. If the cycle $u_1u_2...u_iu_ju_{j+1}...u_m$ is an odd cycle, then this cycle, the cycle $v_1v_2...v_n$ and the path $u_1w_1...w_pv_1$ satisfy the relevant conditions which violates the fact that $m+n+p$ is as small as possible. Therefore this cycle is an even cycle which implies that the cycle $u_iu_{i+1}...u_j$ is an odd cycle, because m is odd. We have either $i > 2$ or $j < m$, since otherwise, the cycle $u_1u_2...u_iu_ju_{j+1}...u_m = u_1u_2u_m$ would be an odd cycle. If $j < m$ then the cycles $u_iu_{i+1}...u_j$, $v_1v_2...v_n$ and the path $u_iu_{i-1}...u_1w_1...w_pv_1$ provide an example with a smaller "$m+n+p$". Similarly, if $i > 2$, then the cycles $u_iu_{i+1}...u_j$, $v_1v_2...v_n$ and the path $u_ju_{j+1}...u_1w_1...w_pv_1$ give an example with a smaller "$m+n+p$".
(3) There are no edges of the form u_iv_j where $i = 1, ..., m$ and $j = 1, ..., n$. This holds because the distance between the cycles $u_1u_2...u_m$ and $v_1v_2...v_n$ is greater than 1.

Now we want to use Proposition 2.2 to show that G is not weakly Farkas. On the contrary, assume that G is weakly Farkas. Furthermore we assume that p is even. The other case, i.e. if p is odd, can be handled similarly. Because of the above properties, we are able to find suitable values $\{a_e\}_{\{e\in G\}}$ such that they satisfy the following

$$a_e = \begin{cases} 1 & e = u_1w_1 \\ -1 & e = u_iu_{i+1} \text{ or } e = v_iv_{i+1} \text{ or } e = w_iw_{i+1} \text{ where } i \text{ is odd} \\ -1 & e = u_mu_1 \text{ or } e = v_nv_1 \\ -1 & e = u_ix \text{ if } x \neq u_{i+1} \text{ and } i \text{ is even} \\ 0 & \text{otherwise} \end{cases}$$

Now consider the rational numbers $\{x_e\}_{\{e\in G\}}$ defined by

$$x_e = \begin{cases} 1 & e = u_1w_1 \text{ or } e = w_pv_1 \\ 1 & e = w_iw_{i+1} \text{ where } i \text{ is even} \\ -1 & e = w_iw_{i+1} \text{ where } i \text{ is odd} \\ 1/2 & e = u_iu_{i+1} \text{ or } e = v_iv_{i+1} \text{ where } i \text{ is even} \\ -1/2 & e = u_iu_{i+1} \text{ or } e = v_iv_{i+1} \text{ where } i \text{ is odd} \\ -1/2 & e = u_mu_1 \text{ or } e = v_nv_1 \\ 0 & \text{otherwise} \end{cases}$$

It is straightforward to check the following

$$\sum_{e\in E(G)} x_e v(e) = 0,$$

$$a_e \leq x_e \leq a_e + 1, \text{for every}\, e \in E(G).$$

Since G is weakly Farkas, by Proposition 2.2, there exist integers

$$a_e \leq y_e \leq a_e + 1 \,\text{where}\, e \in E(G)$$

such that $\sum_{e\in E(G)} y_e v(e) = 0$. Note that it follows from the equality

$$\sum_{e\in E(G)} y_e v(e) = 0$$

that $\sum_{z:xz\in E(G)} y_{xz} = 0$ for every $x \in V(G)$. Since $y_{u_1w_1} \in \{1,2\}$ and $y_{u_1x} \in \{0,1\}$ for every $x \neq w_1, u_2, u_n$, we conclude that $y_{u_1u_2} = -1$ or $y_{u_1u_n} = -1$. Without loss of generality, we may assume that $y_{u_1u_2} = -1$. Since $y_{u_2u_3} \in \{0,1\}$ and $y_{u_2x} \in \{-1,0\}$ for every $x \neq u_3$, it is easy to see that we must have $y_{u_2u_3} = 1$ and $y_{u_2x} = 0$ for every $x \neq u_3, u_1$. Continuing this process, we obtain the following

$$y_e = \begin{cases} 1 & e = u_iu_{i+1} \text{ if } i \text{ is even} \\ -1 & e = u_iu_{i+1} \text{ if } i \text{ is odd} \\ -1 & e = u_mu_1 \\ 0 & e = u_ix \text{ if } 1 < i < m \text{ and } x \notin \{u_{i-1}, u_{i+1}\} \\ 0 & e = u_mx \text{ if } x \notin \{u_{m-1}, u_1\} \end{cases}$$

If there were a vertex $x \notin \{u_2, u_m, w_1\}$ such that $u_1x \in E(G)$ and $y_{u_1x} \neq 0$, then we would have $y_{u_1x} = 1$. Note that $x \notin \{u_1, ..., u_m, w_1, ..., w_p, v_1, ..., v_n\}$ by Properties (1), (2) and (3). But then, we would also have $y_{xu_i} = 0$ if $2 \leq i \leq m$ and $xu_i \in E(G)$. Moreover $y_{xz} \in \{0,1\}$ for every $xz \in E(G)$: It would then follow $\sum_{z:xz\in E(G)} y_{xz} > 0$, a contradiction. So $y_{u_1x} = 0$ if $x \notin \{u_2, u_m, w_1\}$ and $u_1x \in E(G)$. It follows from $\sum_{z:u_1z\in E(G)} y_{u_1z} = 0$ that $y_{u_1w_1} = 2$. Now we have $y_{w_1u_i} = 0$ if $2 \leq i \leq m$ and $w_1u_i \in E(G)$. Moreover, we have $y_{w_1x} \in \{0,1\}$ if $x \neq w_2, u_1$ and $w_1x \in E(G)$. It follows from $\sum_{z:w_1z\in E(G)} y_{w_1z} = 0$ that $y_{w_1w_2} \leq -2$. This contradicts the fact that $y_{w_1w_2} \in \{-1,0\}$. The proof is now complete.

□

References

[1] Aryapoor, M. On linear equations arising in Combinatorics (Part I), arXiv:1406.4466.

[2] Fulkerson, D. R., Hoffman, A. J. and Mcandrew, M. H. Some properties of graphs with multiple edges